

# KIMMTRAK® (tebentafusp-tebn) distribution information

KIMMTRAK is indicated for the treatment of HLA-A\*02:01-positive adult patients with unresectable or metastatic uveal melanoma.<sup>1</sup>

## Dosing and administration<sup>1</sup>

KIMMTRAK is administered once weekly via continuous IV infusion over 15-20 minutes. The starting dose is 20 mcg for week 1, with an increase to 30 mcg for week 2 and 68 mcg for week 3 and subsequent weeks.

## Product information

Each package contains a 100 mcg/0.5 mL single-dose vial of KIMMTRAK, which is a sterile, preservative-free, clear, colorless or slightly yellowish solution.<sup>1</sup>

**10-digit NDC:** 80446-401-01

**11-digit NDC:** 80446-0401-01

**Wholesale acquisition cost (WAC):** \$18,760



### Distributors for physician offices

Specialty distributor	Phone orders	Fax orders and website
Cardinal Health Specialty Pharmaceutical Distribution	877-453-3972 Monday-Friday, 7 AM-6 PM CT (24-hour emergency on call)	<a href="https://specialtyonline.cardinalhealth.com">https://specialtyonline.cardinalhealth.com</a>
McKesson Specialty Health	800-482-6700 Monday-Friday, 7 AM-7 PM CT	Fax: 800-289-9285 <a href="https://mscs.mckesson.com">https://mscs.mckesson.com</a>
Oncology Supply	800-633-7555 Monday-Friday, 8 AM-7 PM CT	<a href="https://www.oncologysupply.com">https://www.oncologysupply.com</a>

### Distributors for hospitals and infusion centers

Specialty distributor	Phone orders	Fax orders and website
ASD Healthcare	800-746-6273 Monday-Thursday, 7 AM-6:30 PM CT; Friday, 7 AM-6 PM CT (24-hour emergency on call)	Fax: 800-547-9413 <a href="https://www.asdhealthcare.com">https://www.asdhealthcare.com</a>
Cardinal Health Specialty Pharmaceutical Distribution	866-677-4844 Monday-Friday, 7 AM-6 PM CT (24-hour emergency on call)	Fax: 614-553-6301 <a href="https://orderexpress.cardinalhealth.com">https://orderexpress.cardinalhealth.com</a>
McKesson Plasma and Biologics	877-625-2566 Monday-Friday, 8 AM-6:30 PM CT	Fax: 888-752-7626 <a href="https://connect.mckesson.com">https://connect.mckesson.com</a>

KIMMTRAK is not currently available through specialty pharmacy orders. HLA-A, human leukocyte antigen-A. An FDA-approved test for the detection of HLA-A\*02:01 genotyping is not currently available.

Please see Important Safety Information including **BOXED WARNING for Cytokine Release Syndrome (CRS)** on next page and see [full Prescribing Information](#).



# Indication and Important Safety Information Including Boxed Warning

## Indication

KIMMTRAK is a bispecific gp100 peptide-HLA-directed CD3 T cell engager indicated for the treatment of HLA-A\*02:01-positive adult patients with unresectable or metastatic uveal melanoma.

## Important Safety Information Including Boxed Warning

### **WARNING: CYTOKINE RELEASE SYNDROME**

**Cytokine Release Syndrome (CRS), which may be serious or life-threatening, occurred in patients receiving KIMMTRAK. Monitor for at least 16 hours following first three infusions and then as clinically indicated.** Manifestations of CRS may include fever, hypotension, hypoxia, chills, nausea, vomiting, rash, elevated transaminases, fatigue, and headache. CRS occurred in 89% of patients who received KIMMTRAK with 0.8% being grade 3 or 4. Ensure immediate access to medications and resuscitative equipment to manage CRS. Ensure patients are euvolemic prior to initiating the infusions. Closely monitor patients for signs or symptoms of CRS following infusions of KIMMTRAK. Monitor fluid status, vital signs, and oxygenation level and provide appropriate therapy. Withhold or discontinue KIMMTRAK depending on persistence and severity of CRS.

### **Skin Reactions**

Skin reactions, including rash, pruritus, and cutaneous edema occurred in 91% of patients treated with KIMMTRAK. Monitor patients for skin reactions. If skin reactions occur, treat with antihistamine and topical or systemic steroids based on persistence and severity of symptoms. Withhold or permanently discontinue KIMMTRAK depending on the severity of skin reactions.

### **Elevated Liver Enzymes**

Elevations in liver enzymes occurred in 65% of patients treated with KIMMTRAK. Monitor alanine aminotransferase (ALT), aspartate aminotransferase (AST), and total blood bilirubin prior to the start of and during treatment with KIMMTRAK. Withhold KIMMTRAK according to severity.

### **Embryo-Fetal Toxicity**

KIMMTRAK may cause fetal harm. Advise pregnant patients of potential risk to the fetus and patients of reproductive potential to use effective contraception during treatment with KIMMTRAK and 1 week after the last dose.

The most common adverse reactions ( $\geq 30\%$ ) in patients who received KIMMTRAK were cytokine release syndrome, rash, pyrexia, pruritus, fatigue, nausea, chills, abdominal pain, edema, hypotension, dry skin, headache, and vomiting. The most common ( $\geq 50\%$ ) laboratory abnormalities were decreased lymphocyte count, increased creatinine, increased glucose, increased AST, increased ALT, decreased hemoglobin, and decreased phosphate.

**Please see [full Prescribing Information](#), including **BOXED WARNING** for CRS.**

**Reference: 1.** Kimmtrak. Package insert. Immunocore Ltd; 2022.

## IMMUNOCORE

Immunocore and KIMMTRAK are registered trademarks of Immunocore Ltd. All other trademarks, company and product names, and logos are the property of their respective owners. ©2022 Immunocore Ltd. All rights reserved. CM-US-TEBE-2100150 January 2022

 **KIMMTRAK**  
(tebentafusp-tebn)  
Injection for Intravenous Use 100 mcg/0.5 mL